



**UNIVERSITI PUTRA MALAYSIA**

**UTILIZATION OF NATURAL FEED FOR GROWTH AND SURVIVAL  
ENHANCEMENT OF *PENAEUS MONODON* JUVENILES IN CULTURE  
SYSTEM AND ITS EFFECTS ON WATER QUALITY**

**FARSHAD SHISHEHCHIAN**

**FSAS 2000 24**

**UTILIZATION OF NATURAL FEED FOR GROWTH AND SURVIVAL  
ENHANCEMENT OF *PENAEUS MONODON* JUVENILES IN CULTURE  
SYSTEM AND ITS EFFECTS ON WATER QUALITY**

**By**

**FARSHAD SHISHEHCHIAN**

**Dissertation Submitted in Fulfillment of the Requirements for the Degree of  
Doctor of Philosophy in the Faculty of Science and Environmental Studies  
Universiti Putra Malaysia**

**November 2000**



**Dedicated to**

**My parents: Esmail Shishehchian and Akram Rahnema**

**My wife: Roya Azimzadeh**

**My daughter & son: Bahar and Hamoon Shishehchian**

**For their Love, Support and Trust**

Abstract of thesis submitted to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

**UTILIZATION OF NATURAL FEED FOR GROWTH AND SURVIVAL  
ENHANCEMENT OF *PENAEUS MONODON* JUVENILES IN CULTURE  
SYSTEM AND ITS EFFECTS ON WATER QUALITY**

By

**FARSHAD SHISHEHCHIAN**

**November 2000**

**Chairperson: Professor Dr. Fatimah Md. Yusoff, Ph.D**

**Faculty: Science and Environmental Studies**

Shrimp feeds on the natural organisms such as plankton, bacteria and benthos. Problems in shrimp farming are due to pollution, diseases and poor feeding practices. There are several advantages in improving natural food in shrimp farming. First, water quality is very much improved since the oxygen evolved in photosynthesis is used in aerobic decomposition of organic matter. Moreover, ammonia in the water are reduced due to uptake by algae. Secondly, the digestibility and acceptability of natural food by shrimp have been well established. Understanding the contribution of natural foods in the shrimp culture system may help to increase the production and reduce the cost of

farming. This study was carried out to demonstrate the importance of natural food in intensive shrimp culture system. Analysis of composition and abundance of macrobenthos in shrimp culture ponds was done by sampling the sediment from six different locations of two shrimp ponds. Gut content analysis was also performed to assess the fullness of shrimp gut. A significant negative correlation ( $r = -0.97$ ,  $p < 0.05$ ) was also observed between the gut content and the abundance of macrobenthos in the pond bottom. There was a decline in the fullness of the shrimp foregut with the reduction of benthic populations in the pond sediments, probably indicating the importance of benthos as food for shrimp in culture system. In another field experiment the effect of bacterial product on development of benthic community in shrimp ponds, was determined. The density of polychaetes and insect-larvae was significantly ( $p < 0.05$ ) higher in ponds with commercial bacteria products than the control, and also showed better feed conversion ratio (FCR). Laboratory experiments showed that the growth rate of *P. monodon* juvenile was significantly ( $p < 0.05$ ) higher when fed with the combination of natural and artificial diets than other treatments. Survival rate was also significantly ( $p < 0.05$ ) higher in those shrimp fed on natural diet compared to the others. The presence of bottom sediment significantly ( $p < 0.05$ ) increased the survival rate of the shrimps. Shrimps fed with artificial diet showed significantly higher excretion of ammonia compare to the others. The results showed that unicellular algae could help shrimp to attain high growth and survival rates. In addition these algae control the water quality parameter such as ammonia-N and nitrite-N. Tanks without shrimp showed significantly lower ( $p < 0.05$ ) rate of nitrogen change compared to the rest of the

treatments, indicating that most of the nitrogenous source was from the shrimp excretion. There was no significant difference among the ingestion rates of three algal species but was relatively high with *Chaetoceros calcitrans* and low with *Tetraselmis tetrahele*. The results indicated that *Penaeus monodon* juvenile was able to ingest and digest the unicellular algae. Feeding with natural feed enhanced the non-specific immune system of the shrimp. The results showed that feeding on chironomid larva significantly ( $p < 0.05$ ) increase the survival rate when shrimp was exposed to the white spot virus (WSV).

The results of this study showed that natural food play an important role in shrimp farming. The presence of suitable natural food may enhance the growth and survival of shrimps in a culture pond and improve the water and sediment quality. This study suggested that the optimum growth of shrimp could be obtained by the right combination of natural and artificial diets.

**PENGUNAAN MAKANAN NATURAL BAGI PENINGKATAN  
PERTUMBUHAN DAN KEMANDIRIAN JUVENIL *PENAEUS MONODON*  
DI DALAM SISTEM KULTUR DAN PENGARUHNYA TERHADAP  
KUALITI AIR**

Oleh

**FARSHAD SHISHEHCHIAN**

**November 2000**

**Pengerusi: Prof. Dr. Fatimah Md. Yusoff**

**Fakulti : Sains dan Pengajian Alam Sekitar**

Udang biasanya memakan makanan yang natural seperti plankton, bakteria dan pelbagai jenis organisma benthik. Masalah yang sering dihadapi oleh industri ternakan udang adalah berpunca daripada pencemaran, penyakit udang yang merebak, dan cara pemberian makanan yang kurang baik. Terdapat beberapa kebaikan untuk meningkatkan kualiti sumber makanan natural dalam kolam udang. Pertama ialah kualiti air dapat dipertingkatkan kerana kandungan oksigen yang dihasilkan daripada proses fotosintesis dapat digunakan dalam penguraian aerobik bahan organik. Tambahan pula, kandungan ammonia dapat dikurangkan oleh proses fotosintesis alga. Di samping itu, pencernaan dan penerimaan makanan semulajadi oleh udang telah diketahui umum. Dengan memahami sumbangan makanan semulajadi dalam perusahaan udang dapat meningkatkan pengeluaran

udang dan mengurangkan kos ternakan. Penyelidikan ini bertujuan untuk menunjukkan kepentingan sumber makanan semulajadi kultur udang intensif. Penganalisisan komposisi dan kepadatan makrobentos dalam kolam kultur udang dilakukan dengan pensampelan sedimen dari enam lokasi dalam dua kolam udang. Analisis kandungan usus juga dilakukan untuk menilai kepenuhan usus udang. Korelasi negatif yang bererti juga dapat diperhatikan di antara kandungan usus dan kepadatan makrobentos di dasar kolam. Terdapat penurunan dalam kepenuhan usus depan udang dengan pengurangan populasi bentos di dalam sedimen kolam, mungkin menunjukkan kepentingan bentos sebagai makanan udang di dalam sistem pengkulturan. Di dalam eksperimen lapangan yang lain, kesan produk bakteria ke atas perkembangan komuniti bentos di dalam kolam udang telah ditentukan. Kepadatan polychaete dan larva serangga adalah lebih tinggi dengan bererti ( $p < 0.05$ ) di dalam kolam yang mengandungi produk bakteria komersial berbanding dengan kawalan, dan juga menunjukkan nisbah penukaran makanan (FCR) yang lebih baik. Eksperimen dalam makmal menunjukkan pertumbuhan *P. monodon* yang jauh lebih tinggi ( $p < 0.05$ ) apabila diberi makanan kombinasi semulajadi dan buatan berbanding dengan rawatan yang lain. Kadar kemandirian udang juga jauh lebih tinggi ( $p < 0.05$ ) untuk udang yang diberi makanan semulajadi jika berbanding dengan yang lain. Kehadiran mendakan telah meningkatkan kadar kemandirian udang dengan bererti ( $p < 0.05$ ). Udang yang diberi makanan buatan menunjukkan perkumuhan ammonia yang jauh lebih tinggi daripada udang yang diberi makanan lain. Keputusan menunjukkan alga unisel boleh membantu udang untuk mencapai kadar pertumbuhan dan kemandirian



yang tinggi. Tambahan pula, alga dapat mengawal parameter kualiti air seperti ammonia-N dan nitrit-N. Rawatan yang tidak mempunyai udang menunjukkan kadar nitrogen yang jauh lebih rendah ( $p < 0.05$ ) berbanding dengan rawatan yang lain. Ini menunjukkan punca nitrogen adalah daripada perkumuhan udang. Keputusan mendapati tidak terdapat perbezaan kadar pengingesan yang bererti bagi ketiga-tiga spesies alga. Walau bagaimanapun, kadar pengingesan *Chaetoceros calcitrans* adalah lebih tinggi dan pencernaan *Tetraselmis tetrahele* adalah lebih rendah. Keputusan menunjukkan juvenil *P. monodon* dapat menginges dan mencerna alga satu sel. Makanan semulajadi juga didapati merangsang sistem imuniti udang yang bukan spesifik. Keputusan menunjukkan bahawa udang yang diberi makan larva kironomid boleh meningkatkan kadar kemandirian udang ( $p < 0.05$ ) apabila udang didedahkan kepada virus bintik putih.

Keputusan penyelidikan ini menunjukkan bahawa sumber makanan semulajadi memainkan peranan yang penting dalam ternakan udang. Kehadiran sumber makanan semulajadi yang sesuai bukan sahaja merangsangkan pertumbuhan dan kemandirian udang dalam kolam, tetapi juga memperbaiki kualiti air dan mendakan kolam. Penyelidikan ini menunjukkan bahawa pertumbuhan udang yang optimum boleh dicapai dengan kombinasi sumber makanan semulajadi dan buatan yang sesuai.

## ACKNOWLEDGEMENTS

I would like to extend my sincere and deep appreciation to Professor Dr. Fatimah Md. Yusoff who provided me with invaluable guidance, real criticism and encouragement throughout the period of the study. I wish to express my heart felt gratitude to Dr. Hishamuddin Omar and Dr. Mohd Salleh Kamarudin, members of my Advisory Committee who helped me by providing important suggestions and some laboratory facilities during the course of my study.

I would to thank Prof. Dr. Tan the Head of the Department of Biology and all staff members for providing the good environment and facilities during my study. My sincere thanks to all staff members and students in Aquatic Research Laboratory, Department of Biology specially to Mr. Azhar Othman, Mrs. Hazel Matias, Mr. Edwardo, Mr. Devaraja, Mr. Hamid Rezai and Mr. Rao.

I would like to extend my most sincere gratitude and thanks to Prof. Dr. Mohamed Shariff for providing me some good suggestions. I also extend my sincere thanks to Miss Abeer for helping with histopathology work. . I would like extend my thanks to Associate Prof. Dr. Jambari H. Ali for helping with identifying the gastropod species.

My special thanks to Miss Alona, Miss Yquen, Miss Shelly, Miss Siti, Miss

Hor Li Phoon and Dr. Paymon.

I would like to appreciate the assistance of Mr. K.C Ting the Managing Director and all staff members of J.W Shrimp Farm, for providing the shrimp culture ponds for some experiments. I also appreciate the assistance of Mr. Jasni and all the staff members of UPM hatchery unit.

My sincere thanks also to the staff members of graduate School for helping me in every possible way. I am grateful to the Malaysian Government for providing funds through Intensification of Research Priority Areas (IRPA Project No. 01-02-04-0165) to carry out this research.

I cannot find appropriate words to express my heartfelt gratitude to my parents for their on-going untiring and guidance since my childhood. I also extend my sincere and deep appreciation to my wife Roya, daughter Bahar and son Hamoon for their moral support and encouragement to me all the time in striving towards this achievement.

I certify that an Examination Committee met on 20<sup>th</sup> November 2000 to conduct the final examination of Farshad Shishehchian on his Doctor of Philosophy thesis entitled "Utilization of Natural Feed for growth and Survival Enhancement of *Penaeus monodon* Juveniles in Culture System and its Effects on Water Quality" in accordance with Universiti Pertanian Malaysia (Higher Degree) Regulations Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Che Roos Saad, Ph.D.  
Institute of Bioscience  
Universiti Putra Malaysia  
(Chairman)

Fatimah Md. Yusoff, Ph.D.  
Professor  
Faculty of Science and Environmental Studies  
Universiti Putra Malaysia  
(Member)

Hishamuddin Omar, Ph.D.  
Faculty of Science and Environmental Studies  
Universiti Putra Malaysia  
(Member)

Mohd. Salleh Kamarudin, Ph.D.  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

  
\_\_\_\_\_  
MOHD GHAZALI MOHAYIDIN, Ph.D.  
Professor / Deputy Dean of Graduate School  
Universiti Putra Malaysia

23 MAR 2001

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy.

---

**MOHD. GHAZALI MOHAYIDIN, Ph.D.**

Professor


Deputy Dean of Graduate School

Universiti Putra Malaysia

Date:

## DECLARATION

I hereby declare that the dissertation is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

  
Farshad Shishehchian

Date: 23/03/2001

## TABLE OF CONTENTS

	Pages
<b>DEDICATION</b> -----	<b>ii</b>
<b>ABSTRACT</b> -----	<b>iii</b>
<b>ABSTRAK</b> -----	<b>vi</b>
<b>ACKNOWLEDGEMENTS</b> -----	<b>IX</b>
<b>APPROVAL SHEETS</b> -----	<b>XI</b>
<b>DECLARATION</b> -----	<b>XIII</b>
<b>LIST OF TABLES</b> -----	<b>XVII</b>
<b>LIST OF FIGURES</b> -----	<b>XX</b>
<b>LIST OF FIGURES</b> -----	<b>XX</b>
<b>LIST OF PLATES</b> -----	<b>XXII</b>
 <b>CHAPTER</b>	
<b>1</b>	
<b>INTRODUCTION</b> -----	<b>1</b>
Background of Study -----	1
Statement of Problem -----	5
Production -----	5
Water and Sediment Quality Problems -----	5
Economic and Management Problems -----	7
Significance of the Study -----	8
Objectives of the study -----	10
 <b>2</b>	
<b>LITERATURE REVIEW</b> -----	<b>11</b>
Biology and Ecology of <i>Penaeus monodon</i> -----	11
Taxonomy -----	11
Morphology -----	11
Distribution -----	12
Morphological Development -----	12
Life history -----	14
Feeding Behavior -----	17
Growth and Energy Reserve in the Shrimp -----	20
Defense System Against Diseases in <i>Penaeus monodon</i> -----	21
Natural Productivity and Shrimp Production -----	24
Stomach Content of Shrimps -----	27
Nutritional Requirements of <i>Penaeus monodon</i> -----	30
Protein and Amino Acid Requirement -----	31
Carbohydrates -----	32
Energy Requirement -----	32
Lipid and Fatty Acid Requirement -----	33
Water and Sediment Quality in <i>Penaeus monodon</i> Culture -----	34
Salinity -----	35

Temperature -----	36
Dissolved oxygen -----	36
pH -----	37
Ammonia, Nitrite, Nitrate and Phosphate -----	37
Excretion and Fecal Matters -----	38
Probiotics in Shrimp Farming -----	39

## 3

<b>GENERAL MATERIALS AND METHODS -----</b>	<b>41</b>
Water Quality -----	41
Total Nitrogen (TN) -----	41
Nitrate Analysis (NO <sub>3</sub> -N) -----	42
Determination of Nitrite (NO <sub>2</sub> -N) -----	43
Determination of Ammonia (Alternative Method) (NH <sub>4</sub> -N) -----	44
Determination of Orthophosphate (PO <sub>4</sub> -P) -----	46
Determination of Total Phosphorus (TP) -----	47
Determination of Hydrogen Sulfide -----	48
Sediment Analysis -----	49
Total Nitrogen -----	49
Total Organic Carbon (Walkey-Black method) -----	50
Total Available Phosphorus -----	51
Proximate Composition -----	52
Protein -----	52
Lipid -----	53
Fatty Acid Determination (Direct Methods) -----	54
Soluble Carbohydrate -----	55
Estimation of Chlorophyll a -----	56
Rotifer and Microalgae Culture -----	57
Rotifer Culture -----	57
Culture of Microalgae -----	58

## 4

<b>MACROBENTHOS AND FEEDING BEHAVIOR OF SHRIMP IN CULTURE PONDS -----</b>	<b>61</b>
Introduction -----	61
Macrobenthos in shrimp pond -----	62
Effects of commercial bacterial products on benthic community -----	63
Materials and Methods -----	66
Experiment I: Macro-benthos in the <i>Penaeus monodon</i> Culture Systems -----	66
Experiment II: Effect of Commercial Bacterial Products on the Benthic Community -----	67
Results and Discussion -----	71
Experiment I -----	71
Experiment II -----	86



## 5

<b>GROWTH AND SURVIVAL RATE OF <i>PENAEUS MONODON</i> IN DIFFERENT CULTURE CONDITIONS</b>	<b>98</b>
Introduction	98
Materials and Methods	102
Experiment I: Growth, Survival and Biochemical Composition of <i>Penaeus monodon</i> Juvenile Fed with Natural and Artificial Diets	102
Experiment II: Growth and Survival of <i>Penaeus monodon</i> Juvenile Fed with Different Diets and Culture in Tanks with the Presence and Absence of Bottom Sediment	106
Experiment III: Effect of natural diets on resistance to white spot syndrome virus (WSSV) in <i>Penaeus monodon</i> .	108
Results and Discussion	110
Experiment II	123
Experiment III	129

## 6

<b>EFFECTS OF DIFFERENT DIETS ON THE NITROGENOUS EXCRETION OF <i>PENAEUS MONODON</i></b>	<b>135</b>
Introduction	135
Materials and Methods	138
Experiment I: Nitrogenous Excretion of <i>Penaeus monodon</i> Post Larvae Fed with Natural and Artificial Diets	138
Experiment II: The Nitrogenous Contents of Water in Tanks with Shrimps Fed on Different Algal Species	140
Results and Discussion	142
Experiment I	142
Experiment II	148

## 7

<b>INGESTION RATE OF <i>PENAEUS MONODON</i> JUVENILE FED WITH <i>CHAETOCEROS CALCITRANS</i>, <i>TETRASELMIS TETRAHELE</i> AND <i>NANNOCHLOROPSIS OCULATA</i></b>	<b>152</b>
Introduction	152
Material and Methods	153
Results and Discussion	155

## 8

<b>SUMMARY</b>	<b>162</b>
<b>REFERENCES</b>	<b>169</b>
<b>VITAE</b>	<b>190</b>

## LIST OF TABLES

Table 1-1: World farmed shrimp production in 1995 (Hagler, 1997).....	2
Table 2-1: Some basic information about the most important species of <i>Penaeidae</i> (Jung and Co, 1988). ....	15
Table 2-2: Life history of the <i>Penaeus monodon</i> (Motoh, 1981).....	19
Table 2-3: Animals that are commonly observed in Hawaiian prawn pond .....	25
Table 2-4: Morphological parameters of <i>Penaeus monodon</i> fed with different types of algae and pellet (Primavera and Gacutan, 1989).....	26
Table 2-5: Protein requirements of <i>Penaeus monodon</i> .....	31
Table 2-6: Recommended water parameters for successful culture of <i>Penaeus monodon</i> (Fast and Lester, 1992; Dall et al., 1990). ....	36
Table 3-1: Composition of Conwy media (Tompkins et al., 1995).....	60
Table 4-1: List of macrobenthos found on Carey Island shrimp culture ponds during 120-culture cycle (September to December 1996).....	72
Table 4-2: Mean percentage ( $\pm$ standard error) of total nitrogen, phosphorus and organic matter in sediments of different treatment ponds for the whole culture period. Means in a row with different letter are significantly different ( $P < 0.05$ ). ....	86
Table 4-3: List of the major macrobenthos found in shrimp culture ponds during the culture cycle in all treatment ponds.....	89
Table 4-4: Mean ( $\pm$ standard error) density of the benthos during the culture period in different treatments. Means ( $\pm$ standard error) in a column without a letter in common is significantly different ( $P < 0.05$ ). ....	90
Table 4-5: Mean ( $\pm$ standard error) density of the edible benthos (Polychaeta and insect larvae) during the culture period in different treatments and different time. Means ( $\pm$ standard error) in a row without a letter in common is significantly different ( $P < 0.05$ ). ....	91
Table 4-6. Mean value ( $\pm$ Standard error) of water parameters in different treatments and control.....	93
Table 4-7: The most abundance items, which were identified in the foreguts of shrimp in different treatments and control. Values the percentages forgut over all examined stomach which particular food items were found (n=9). ....	97
Table 4-8: Mean ( $\pm$ standard error) of total shrimp production and feed conversion ratio (FCR) in each treatment and control. Means in a row without a common letter are significantly different ( $P < 0.05$ ). ....	97

Table 5-1: Final weight (mg), weight gain <sup>1</sup> (%), final length (mm), survival (%) and SGR <sup>2</sup> (%) of <i>Penaeus monodon</i> . Values are mean ( $\pm$ standard error). Means in a row without a common letter common are significantly different ( $P < 0.05$ , $n$ =survived shrimps).....	113
Table 5-2: Proximate composition (%) of <i>Penaeus monodon</i> during the trial (mean $\pm$ standard error).....	116
Table 5-3: Proximate composition (%) of different diets, which used during the trial (mean $\pm$ standard error, $n = 3$ ).....	116
Table 5-4: Fatty acid composition of shrimp in different treatments (mean $\pm$ standard error).....	118
Table 5-5: Fatty acid composition of different diet (mean $\pm$ standard error).....	119
Table 5-6: Dissolved oxygen (mg/L), temperature ( $^{\circ}$ C), NH <sub>4</sub> -N (mg/L) and pH during the experiment II (mean $\pm$ standard error) ( $P < 0.05$ , $n=9$ ).....	123
Table 5-7: Final weight (mg), weight gain <sup>1</sup> (%), final length (mm), survival (%) and SGR <sup>2</sup> (%) of <i>Penaeus monodon</i> . Values are mean ( $\pm$ standard error). Means in a row without a letter in common are significantly different ( $P < 0.05$ , $n$ =survived shrimps). ....	126
Table 5-8: Proximate composition (%) of <i>Penaeus monodon</i> during the trial. (mean $\pm$ standard error). Means in a row without a letter in common are significantly different ( $P < 0.05$ , $n=9$ ). ....	127
Table 5-9: Fatty acid composition (%) of shrimp in different treatments (mean $\pm$ standard error).....	128
Table 5-10: Proximate composition (%) of different diets, used during the trial (mean $\pm$ standard error, $n = 3$ ).....	129
Table 5-11: Survival rate of <i>Penaeus monodon</i> , exposed to white spot syndrome virus (WSSV) in different treatments (mean $\pm$ standard error).....	130
Table 5-12: Results of two way ANOVA. Means in a row for each dietary treatment and control without a letter in common are significantly different ( $P < 0.05$ ). ....	131
Table 6-1: The mean of pH values in different treatments.....	142
Table 6-2: The mean rate of concentration change (mg/L/hr) of total NH <sub>4</sub> -N and NO <sub>2</sub> -N in different treatments. Means in same column having similar superscript are not significantly different ( $p > 0.05$ ) (mean $\pm$ standard error).....	144
Table 6-3: The mean concentration (mg/L) of total NH <sub>4</sub> -N in different control .....	148
Table 6-4: The mean rate of concentration change (mg/hr) of total NH <sub>4</sub> -N and NO <sub>2</sub> -N among different treatment those in same column	

having similar superscript are not significantly different ( $P>0.01$ ). .....	149
Table 7-1: Dissolved oxygen (mg/L), temperature ( $^{\circ}\text{C}$ ), pH, total ammonia-N (mg/L) and $\text{PO}_4\text{-P}$ (mg/L) during the trial in the experimental tanks (means $\pm$ standard error). Means in a row without a letter in common are significantly different ( $P<$ 0.05).....	156
Table 7-2: Dissolved oxygen (mg/L), temperature ( $^{\circ}\text{C}$ ), pH, total ammonia-N (mg/L) and $\text{PO}_4\text{-P}$ (mg/L) during the trial in the control tanks (means $\pm$ standard error). No significant difference ( $p>0.05$ ) was detected between means of treatments.....	157
Table 7-3: Ingestion rate (mg/ mg dry weight shrimp/hr) and clearing rate among the different treatment. Means ( $\pm$ standard error) in a row without a letter in common are significantly different ( $P< 0.05$ ). .....	157

## LIST OF FIGURES

Figure 2-1: Various parts of juvenile <i>Penaeus monodon</i> (Motoh, 1981).....	13
Figure 2-2: Diagram of the life history of the <i>Penaeus monodon</i> (Motoh, 1981) .....	16
Figure 4-1: Location of sediment sampling sites and typical water circulation in ponds. LE (left effluent), LI (left influent), ME (mid effluent), MM (mid mid), MI (mid influent), RM (right mid) are the sampling sites. ..	67
Figure 4-2: Location of sediment sampling sites and typical water circulation in ponds.....	70
Figure 4-3: Fluctuations of (a) temperature and salinity, and (b) dissolved oxygen and pH during the culture cycle. The values represent the mean of two replicates. ....	75
Figure 4-4: Percentage of major taxonomic groups of macrobenthos in the sediments during the entire culture period. ....	76
Figure 4-5: Changes in the density of macrobenthos in sediments at 30-day intervals after stocking.....	77
Figure 4-6: Percentage of distribution for each individual taxon at various sampling time.....	78
Figure 4-7: Relationship between density of macrobenthos and fullness of gut of shrimp. ....	79
Figure 4-8: Fluctuation of the percentage of total nitrogen in the sediment during the entire culture period (mean $\pm$ standard error). ....	87
Figure 4-9: Percentages of total polychaetes and insect larvae (edible benthos) in the different treatments during the whole culture cycle. ....	88
Figure 4-10: Percentages of total polychaetes and insect larvae in different treatments at different phases of the culture period. ....	92
Figure 4-11: Percentage of the density of each individual taxon in different treatments and control at various sampling times. ....	94
Figure 4-12: Percentage of the fullness of shrimp foregut at different time during 24-hr study.....	95
Figure 5-1: Correlation ( $r^2 > 0.98$ ) between number and weight of <i>Chaetoceros</i> <i>calcitrans</i> (a), <i>Tetraselmis</i> sp. (b) and <i>Nannochloropsis oculata</i> (c). ....	112
Figure 6-1: Change of pH in different treatments during 103 hours.....	143
Figure 6-2: Rates of concentration change of $\text{NH}_4\text{-N}$ (a) and $\text{NO}_2\text{-N}$ (b) in different treatments during 103 hours.....	147
Figure 6-3: Rates of concentration change of $\text{NH}_4\text{-N}$ (a) and $\text{NO}_2\text{-N}$ (b) in different treatments during 118 hours. Bars indicate maximum and minimum values. ....	151
Figure 7-1: The changes in the density (mg) of the algae in control tanks, <i>Chaetoceros calcitrans</i> , <i>Tetraselmis tetrahele</i> and <i>Nannochloropsis</i> <i>oculata</i> .....	159

<i>Figure 7-2: The changes in the concentration (mg) of the algae during the experiment, of <i>Tetraselmis tetrahele</i>, <i>Chaetoceros calcitrans</i> and <i>Nannochloropsis oculata</i>.</i> .....	160
<i>Figure 7-3: The changes in the concentration (mg/L) of NH<sub>4</sub>-N and PO<sub>4</sub>-P in the experiment and control tanks during the experiment.</i> .....	161

## LIST OF PLATES

Plate 3-1: Mass culture of the different algae species by using the plastic bag .....	59
Plate 4-1: Different types of gastropod found during the culture period (A1, C1 and D1: Littorinidae; B2: Stenothyridae; C3: <i>Melaroides tulerculata</i> ).....	80
Plate 4-2: Different types of oligochaetes(a) polychaetes (b) and insect larvae (c), found during the culture period.....	81
Plate 4-3: One species of crustacean (Macrura), found during the culture period. ....	82
Plate 4-4: Different types of bivalve found during the culture period (1and 3: Veneroida).....	83
Plate 4-5: The appearance of (a) full stomach, (b) empty stomach and (c) opened stomach of shrimp. ....	84
Plate 4-6: Stomach contents of shrimp. (a) Remains of copepod (b) remains of plant material, (b and c) artificial diet and sand, (d) remain of gastropod (d).....	85
Plate 5.1: Different histopatological results of the shrimp tissue. ....	134



## CHAPTER 1

### INTRODUCTION

#### Background of Study

One of the most important cultured organisms is the marine shrimp. The history of marine shrimp culture is not well documented. Its origin traces to the earliest brackishwater and marine ponds. These primary ponds, which date back thousands of years in the Mediterranean area, and to the 15<sup>th</sup> century AD in Indonesia, were typically stocked with fish, shrimp, and other biota through passive flooding (Fast and Lester, 1992). Desirable seed, as well as unwanted predators and competitors were all carried into the pond with source waters where they grew for some time, and were then harvested. This culture method, which is called “catch and hold” culture, is still commonly used today. Where land and labor costs are low, and pond improvement costs have been largely depreciated, this approach can still be profitable with low risk (Fast and Lester, 1992).

Modern shrimp farming was born in the 1930s when Motosaku Fujinaga, a graduate of University of Tokyo, succeeded in spawning the kuruma shrimp (*Penaeus japonicus*). He cultured larvae through to market size in the laboratory and succeeded in mass-producing them on a commercial scale. For more than 40 years, he generously shared his findings and published papers in 1935, 1941,



1942 and 1967. Emperor Hirohito honored him with the title “Father of Inland Japonicus Farming” (Shigueno, 1975).

Worldwide, the majority of shrimp culture takes place in extensive, semi-intensive and intensive earthen ponds. The tiger prawn, *Penaeus monodon* (Fabricius), is the most extensively cultured crustacean in the South-East Asia. This is probably due to high growth rate capacity for adaptation to various culture system (Alava and Lim, 1983) and good response to commercial feed (Lee, 1971).

Table 1-1: World farmed shrimp production in 1995 (Hagler, 1997).

	Percent of production	Heads-on production (t)	Hectares in production	Number of farms
<b>Thailand</b>	31	220,000	90,000	20,000
<b>Ecuador</b>	14	100,000	125,000	1,900
<b>Indonesia</b>	11	80,000	300,000	4,000
<b>China</b>	10	70,000	13,000	6,000
<b>India</b>	8	60,000	80,000	5,000
<b>Vietnam</b>	7	50,000	225,000	2,000
<b>Bangladesh</b>	4	30,000	130,000	9,000
<b>Mexico</b>	2	12,000	14,000	250
<b>Columbia</b>	2	11,000	3,000	30
<b>Honduras</b>	1	10,000	11,000	36
<b>Other</b>	11	78,000	30,670	2,152
<b>TOTAL</b>	100	721,000	1,021,670	50,368

According to data received from the United Nations Food and Agriculture Organization (FAO), world shrimp production set a new record high volume in